

Lifetime and baseline alcohol intakes and risk of pancreatic cancer in the European Prospective Investigation into Cancer and Nutrition study

Ref: IJC-17-2412

Sabine Naudin¹, Kuanrong Li¹, Tristan Jaouen¹, Nada Assi¹, Cecilie Kyrø², Anne Tjønneland², Kim Overvad³, Marie-Christine Boutron-Ruault^{4,5}, Vinciane Rebours⁶, Anne-Laure Védie⁶, Heiner Boeing⁷, Rudolf Kaaks⁸, Verena Katzke⁸, Christina Bamia^{9,10}, Androniki Naska^{9,10}, Antonia Trichopoulou^{9,10}, Franco Berrino¹¹, Giovanna Tagliabue¹², Domenico Palli¹³, Salvatore Panico¹⁴, Rosario Tumino¹⁵, Carlotta Sacerdote¹⁶, Petra H. Peeters^{17,18}, Bas Bueno-de-Mesquita^{18,19,20}, Elisabete Weiderpass^{21,22,23,24}, Inger Torhild Gram²¹, Guri Skeie²¹, Maria-Dolores Chirlaque^{25,26,27}, Miguel Rodríguez-Barranco^{26,28}, Aurelio Barricarte^{26,29,30}, J.Ramón Quirós³¹, Miren Dorronsoro³², Ingegerd Johansson³³, Malin Sund³⁴, Hanna Sternby³⁵, Kathryn E Bradbury³⁶, Nick Wareham³⁷, Elio Riboli³⁸, Marc Gunter³⁹, Paul Brennan⁴⁰, Eric J Duell⁴¹, Pietro Ferrari^{1*}

¹ Nutritional Methodology and Biostatistics group, International Agency for Research on Cancer, Lyon, France; ² Danish Cancer Society Research Center, Copenhagen, Denmark; ³ Section for Epidemiology, Department of Public Health, Aarhus University, Aarhus, Denmark; ⁴ CESP, INSERM U1018, University of Paris-Sud, UVSQ, University of Paris-Saclay, Villejuif, France; ⁵ Institut Gustave Roussy, Villejuif, France; ⁶ Pancreatology Unit, Beaujon Hospital, Clichy, France, INSERM U1149, University Paris 7, Paris, France; ⁷ Department of Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke (DIfE), Potsdam, Germany; ⁸ Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany; ⁹ Hellenic Health Foundation, Athens, Greece; ¹⁰ WHO Collaborating Center for Nutrition and Health, Unit of Nutritional Epidemiology and Nutrition in Public Health, Department of Hygiene, Epidemiology and Medical Statistics, School of Medicine, National and Kapodistrian University of Athens, Greece; ¹¹ Department of Preventive & Predictive Medicine, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy; ¹² Lombardy Cancer Registry Unit, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy; ¹³ Cancer Risk Factors and Life-Style Epidemiology Unit, Cancer Research and Prevention Institute (ISPO), Florence, Italy; ¹⁴ Department of Clinical and Experimental Medicine, University Federico II, Naples, Italy; ¹⁵ Cancer Registry and Histopathology Department, Civic M.P.Arezzo Hospital, Ragusa, Italy, Ragusa, Italy; ¹⁶ Unit of Cancer Epidemiology, Città della Salute e della Scienza University, Hospital and Center for Cancer Prevention (CPO), Turin, Italy; ¹⁷ Department of Epidemiology, Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, The Netherlands; ¹⁸ Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, United Kingdom; ¹⁹ Department for Determinants of Chronic Diseases (DCD), National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands; ²⁰ Department of Social & Preventive Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia; ²¹ Department of Community

1 Medicine, Faculty of Health Sciences, University of Tromsø, The Arctic University of
2 Norway, Tromsø, Norway; ²² Department of Research, Cancer Registry of Norway, Institute
3 of Population-Based Cancer Research, Oslo, Norway; ²³ Department of Medical
4 Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden; ²⁴ Genetic
5 Epidemiology Group, Folkhälsan Research Center, Helsinki, Finland; ²⁵ Department of
6 Epidemiology, Regional Health Council, IMIB-Arrixaca, Murcia, Spain; ²⁶ CIBER of
7 Epidemiology and Public Health (CIBERESP), Madrid, Spain; ²⁷ Department of Health and
8 Social Sciences, University of Murcia, Murcia, Spain; ²⁸ Biosanitary Investigation Institute
9 (IBS) of Granada, University Hospital and University of Granada, Granada, Spain; ²⁹ Navarra
10 Public Health Institute, Pamplona, Spain; ³⁰ Navarra Institute for Health Research (IdiSNA),
11 Pamplona, Spain; ³¹ Public Health Directorate, Asturias, Spain; ³² Subdirección de Salud
12 Pública de Gipuzkoa, Gobierno Vasco, San Sebastian, Spain; ³³ Department of Odontology,
13 Cariology, Umeå University, Umeå, Sweden; ³⁴ Department of Surgical and Perioperative
14 Sciences, Umeå University, Umeå, Sweden; ³⁵ Department of Surgery, Institution of Clinical
15 Sciences Malmö, Lund University, Malmö, Sweden; ³⁶ Cancer Epidemiology Unit, Nuffield
16 Department of Population Health, University of Oxford, Oxford, United Kingdom; ³⁷ MRC
17 Epidemiology Unit, University of Cambridge School of Clinical Medicine, Cambridge
18 Biomedical Campus, Cambridge, United Kingdom; ³⁸ School of Public Health, Imperial
19 College London, United Kingdom; ³⁹ Nutrition and Epidemiology group, International
20 Agency for Research on Cancer, Lyon, France; ⁴⁰ Genetic Epidemiology group, International
21 Agency for Research on Cancer, Lyon, France; ⁴¹ Unit of Nutrition and Cancer, Catalan
22 Institute of Oncology (ICO-Idibell), Barcelona, Spain;

24 **Corresponding Author**

25 Pietro Ferrari, PhD
26 Nutritional Methodology and Biostatistics Group
27 International Agency for Research on Cancer, WHO
28 150 cours Albert Thomas
29 69372 Lyon CEDEX 08, France
30 Tel. +33 472 738 031
31 E-mail: ferrarip@iarc.fr

33 **Abbreviations**

34 EPIC: European prospective investigation into cancer and nutrition
35 HR: hazard ratio
36 CI: confidence interval
37 IARC: International Agency for Research on Cancer
38 WCRF/AICR: World cancer research fund / American institute of cancer research

40 **Article category**

41 Cancer epidemiology

Novelty and Impact (Words=70)

Pancreatic cancer (PC) has been inconsistently associated with alcohol consumption owing to the challenge of investigating a rare disease in prospective studies, providing a limited number of incident events. Through a comprehensive evaluation that included 1,283 incident cases, our study indicated that baseline and lifetime alcohol intakes were positively related to PC, with stronger risks estimated for beer and spirits/liquors than wine intake. Associations were not modulated by smoking habits.

Abstract (Words=243)

Recent evidence suggested a weak relationship between alcohol consumption and pancreatic cancer (PC) risk. In this study, the association between lifetime and baseline alcohol intakes and the risk of PC was evaluated, including the type of alcoholic beverages and potential interaction with smoking. Within the European Prospective Investigation into Cancer and Nutrition (EPIC) study, 1,283 incident PC (57% women) were diagnosed from 476,106 cancer-free participants, followed up for 14 years. Amounts of lifetime and baseline alcohol were estimated through lifestyle and dietary questionnaires, respectively. Cox proportional hazard models with age as primary time variable were used to estimate PC hazard ratios (HR) and their 95% confidence interval (CI). Alcohol intake was positively associated with PC risk in men. Associations were mainly driven by extreme alcohol levels, with HRs comparing heavy drinkers (>60 g/day) to the reference category (0.1-4.9 g/day) equal to 1.77 (95% CI: 1.06, 2.95) and 1.63 (95% CI: 1.16, 2.29) for lifetime and baseline alcohol, respectively. Baseline alcohol intakes from beer (>40 g/day) and spirits/liquors (>10 g/day) showed HRs equal to 1.58 (95% CI: 1.07, 2.34) and 1.41 (95% CI: 1.03, 1.94), respectively, compared to the reference category (0.1-2.9 g/day). In women, HR estimates did not reach statistical significance. The alcohol and PC risk association was not modified by smoking status. Findings from a large prospective study suggest that baseline and lifetime alcohol intakes were positively associated with PC risk, with more apparent risk estimates for beer and spirits/liquors than wine intake.

1 **Introduction**

2 Pancreatic cancer (PC) is a major public health concern. It is one of the most fatal cancers
3 worldwide, accounting for a mortality-incidence ratio close to 1, and a 7% survival beyond
4 five years after diagnosis.^{1,2} The total number of deaths due to PC is expected to rise in the
5 coming years among the American and European populations and is set to surpass breast,
6 prostate, and colorectal cancers to become the second leading cause of cancer-related death
7 after lung cancers.^{3,4} This evidence highlights the importance of understanding risk factors of
8 PC to enhance its primary prevention.

9 The majority of PC cases currently occurs in high-income countries, such as the United States
10 and Western European countries, where incidence rates are nearly three times higher than in
11 middle- and low-income countries.⁵ This incidence pattern suggests that PC occurrence is
12 related to lifestyle factors specifically prevalent in the Western world. The etiology of PC has
13 been extensively researched, leading to the identification of tobacco smoking, obesity, type-II
14 diabetes mellitus and chronic pancreatitis as well as inherited genetic disorders as major risk
15 factors.⁶⁻⁹

16 In 2012, international expert panels reviewed the association between alcohol and cancer and
17 considered the epidemiologic evidence for PC inconsistent, highlighting the possibility of
18 residual confounding by smoking and the lack of knowledge on whether results differ by type
19 of alcoholic beverages.^{6,10} The most recent prospective studies suggested that alcohol
20 consumption may increase PC risk but with an excess risk limited to high levels of
21 consumption.¹¹⁻¹⁴ The majority of these investigations primarily focused on baseline alcohol
22 intake, whereas two early analysis from the European Prospective Investigation on Cancer
23 and Nutrition (EPIC) study indicated that neither baseline nor cumulative lifetime alcohol
24 intake were related to PC risk.^{15,16} Recent meta-analyses have shown that alcohol intake
25 increased the risk of PC by at least 15% in heavy drinkers consuming more than 25 g/day
26 when compared to light drinkers.^{17,18} Although the association was also investigated among
27 never smokers, as well as the interaction with tobacco smoking,^{11,12,14} it has been more often
28 explored in case-control studies in comparison to prospective studies¹⁹ due to the small
29 number of cases being both heavy drinker and never smoker.

30 In the light of these findings, relationship between alcohol intake and PC risk was
31 comprehensively examined in the EPIC study involving a larger number of incident PC cases
32 than earlier evaluations,^{15,16} and presenting risk estimates according to lifetime and baseline
33 intakes, as well as according to the type of alcoholic beverages and smoking habits.

1 **Material and Methods**

2 EPIC is an ongoing multicenter prospective study aiming to investigate prospectively the
3 etiology of cancer in relation to diet, lifestyle and environmental factors, and for which the
4 study design has been previously describe in detail.²⁰ From 1992 to 2000, a total of 521,324
5 participants were recruited across 10 European countries, mostly from the general population,
6 of which 70% are women, aged from 35 to 70 years. Exceptions were the French cohort
7 (members of a health insurance for school and university employees), some of the Spanish
8 and Italian centers (blood donors), Utrecht and Florence sub-cohorts (only breast cancer
9 screening participants), and Oxford sub-cohort (vegetarians and ‘health conscious’
10 participants). The cohorts of France and Norway and the national sub-cohorts of Utrecht and
11 Naples consist of women only. Approval for this study was obtained from the relevant ethical
12 review boards of the participating institutions and study participants provided informed
13 consent before they completed diet, lifestyle and medical questionnaires at baseline.

14 *Assessment of alcohol intake and covariates.* Diet was assessed at recruitment by validated
15 center-/country- specific dietary questionnaires²⁰ designed to capture local-dietary habits with
16 high compliance.²¹ Data on weight and height (self-reported in France, Norway and the UK
17 Oxford center), occupational and physical activities, previous illness, smoking status and
18 lifetime alcohol intake were collected through lifestyle questionnaires.

19 Baseline alcohol intake was computed from the number of glasses of beer and/or cider, wine,
20 sweet liquors and/or distilled spirits, and fortified wines drunk per day or week during the 12
21 months preceding recruitment. For each country, an average daily alcohol intake expressed in
22 grams per day was calculated based on the standard glass volume and ethanol content for each
23 type of alcoholic beverage using information collected through 24-hour dietary recalls from a
24 subgroup of the cohort.^{22–24}

25 Lifetime alcohol consumption was measured through the number of glasses from the different
26 types of beverages consumed per week at 20, 30, 40, and 50 years of age, including the intake
27 at recruitment. The average lifetime alcohol intake was calculated as a weighted average of
28 intakes at different ages with weights equal to the time of exposure to alcohol at different
29 ages. Information was available for 76.3% of the study participants, as data on lifetime
30 alcohol exposure were not collected in Naples (Italy), Bilthoven (The Netherlands), Sweden,
31 and Norway.

32 *Ascertainment of disease outcome and vital status.* The identification of cancer cases during
33 follow-up was based on population cancer registries in 7 of the participating countries

(Denmark, Italy, Netherlands, Norway, Spain, Sweden, and the United Kingdom), and on a combination of methods, including health insurance records, contacts with cancer and pathology registries, and active follow-up of EPIC participants and their next of kin (France, Germany, and Greece). Mortality data were collected from, either the cancer, or mortality registries at the regional or national level. Currently, the vital status is known for 98.4% of all EPIC participants, as well as the proportion of participants who had emigrated to another country, withdrew or had unknown vital status (1.6%).

For the present study, we used information on the most recent vital status and cancer diagnosis update. The follow-up period ended as follows: December 2009 (Varese, Murcia), December 2010 (Florence, Ragusa, Turin, Asturias, Bilthoven, and Utrecht), December 2011 (Granada, Navarra, San Sebastian, and Cambridge), December 2012 (Oxford, Umeå, Denmark, and Norway), and December 2013 (Malmö). For France, Germany, Greece and Naples, the end of follow-up was considered to be the last known contact with study participants: June 2008 for France, December 2009 for Heidelberg and Potsdam, December 2010 for Naples and December 2012 for Greece. Cases of PC defined in this study were primary incident exocrine tumor of the pancreas. They were coded according to International Classification of Diseases-Oncology (3rd edition), including all invasive pancreatic cancers coded as C25 (C25.0–C25.3, C25.7–C25.9). As they represent around 95% of PC cases, this study focused only on exocrine PC, while endocrine tumors of the pancreas were not considered (C25.4). Microscopically confirmed PC represented 67% of the cases (n=854) based on histology, cytology or hematology reports. Other cases were obtained from clinical or surgical observations (n=344), medical imaging techniques (n=57), death certificates (n=17) and laboratory techniques (n=11).

Statistical analysis. EPIC participants without lifestyle or dietary information (n= 6,902), participants with ratio of estimated energy intake over energy requirement in the top or bottom 1% (n=10,241),²⁵ prevalent cancer cases (n=21,401), PC cases with missing date of diagnosis (n=18), participants with missing follow-up information (n=18) and PC cases having a neuroendocrine or endocrine tumor (n=54) were excluded. For lifetime alcohol analysis, participants without information on past alcohol use were excluded (n=112,841).

The association between alcohol intake and PC incidence was evaluated using multivariable Cox proportional hazard models. Age was the primary time variable, and Breslow's method was adopted for handling ties.²⁶ The time at entry was the age at recruitment, whereas the exit time was the age at cancer diagnosis, death, loss, or end of follow-up, whichever came first.

All models were stratified by study center to control for different effects in questionnaires, follow-up procedures and other center-specific features.²⁵ To further control for the effect of age as possible confounding, models were also stratified by age at recruitment in 1-year categories. Separate models were run by gender to account for the behavioral differences of alcohol uses between men and women. Baseline and lifetime alcohol intake were first modeled by categories, as non-consumers, 0.1-4.9 g/day (reference category), 5-14.9 g/day, 15-29.9 g/day, 30-59.9 g/day and >60 g/day. In women, the last two categories were collapsed into a ≥ 30 g/day group. In analyses on lifetime alcohol intake, former drinkers at baseline were separated out from never consumers. Overall tests for significance of HRs related to alcohol in categories were determined by p-values comparing Wald test statistics to a χ^2 distribution with degree of freedom equal to the number of alcohol categories minus one. Analyses were also carried out in continuous, expressing HRs per 12 g/day increase of alcohol intake as 12 grams of alcohol corresponds to about one standard glass of either wine, beer or spirits/liquors. Tests for trend were computed accordingly.

The following confounding variables were consistently included in all analyses: smoking intensity (never; current, 1-15 cig/day; current, 16-25 cig/day; current ≥ 26 cig/day; former, quit <10 years; former, quit 11-20 years; former, quit ≥ 20 years, current, pipe/cigar occasionally, unknown (n=7,921)), education level (no degree, primary school, secondary school, technical or professional school, university degree, unknown (n=10,706)), physical activity index (inactive, moderately inactive, moderately active, active, unknown (n=8,823))²⁷, type 1 and type 2 diabetes mellitus status combined (no, yes, unknown (n=2,324)), body mass index (BMI) in kg.m^{-2} (continuous), height in cm (continuous). The inclusion of energy intake from non-alcohol sources to perform iso-caloric comparisons and partially control for errors in alcohol estimation did not alter the magnitude or risk estimates, and was not pursued. Models evaluating lifetime alcohol consumption were further adjusted on the duration of alcohol drinking (in years), time since quitting (in years), and an indicator variable for drinkers. Associations between alcohol subtypes, namely beer, wine and spirits/liquors and PC were assessed in adjusted models for energy intake from alcohol sources other than the one under evaluation using the following categories: never, 0.1–2.9 g/day (reference), 3–9.9, 10–19.9, 20–39.9 and ≥ 40 g/day. For women, the two last categories were merged into a ≥ 20 g/day group. All models were compatible with the proportional hazards assumption, assessed through analyses of Schoenfeld residuals.²⁸

Dose-response analyses were performed for baseline and lifetime alcohol intake in men. Potential departures from linearity in the association between alcohol intakes and PC were examined by fitting restricted cubic spline models²⁹ with alcohol category-specific knots placed at 0.1, 5, 30, 60 and 100. Non-linearity was evaluated by comparing the difference in log-likelihood of models with linear term and fractional polynomials to a χ^2 distribution.

Effect modification in the relationship between alcohol and PC risk by, in turn, smoking status (never, current smokers), sex and country was evaluated through comparisons of models with and without interaction terms. The differences in log-likelihood were compared to a χ^2 distribution, with degrees of freedom equal to the total number of interaction terms minus one. For analysis by smoking status, parameter estimates were not altered by the inclusion in the models of smoking duration and age at smoking initiation (data not shown).

Sensitivity analyses were performed to assess the robustness of the findings. First, as reverse causation may bias the association between alcohol and PC, cases occurring during the first 2 years of follow-up were further excluded. Second, models on baseline alcohol intake in women were further adjusted for baseline information for menopausal status, ever use of hormone therapy, and number of full-term pregnancies. Finally, in the absence of information on chronic pancreatitis in EPIC, a sensitivity analysis was carried out to account for the potential confounding role of chronic pancreatitis (Z) between baseline alcohol intake (X) and risk of pancreatic cancer (D) using external information.³⁰ A PC HR for baseline heavy drinkers (>60g/day) vs. moderate drinkers (0.1-4.9g/day) not adjusted for chronic pancreatitis in EPIC was estimated as large as 1.64 (95% CI: 1.22, 2.21), for men and women combined. Assuming values from the literature for relative risk estimates of chronic pancreatitis associated with alcohol intake greater than 25 g/day compared to the never drinkers ranging from 2 to 6,^{31,32} pancreatitis prevalence among moderate drinkers ranging from 0.005 to 0.02³² and relative risk estimates of PC associated with chronic pancreatitis ranging from 1.5 to 15,³³⁻³⁵ PC HR for heavy drinkers vs. moderate drinkers adjusted for chronic pancreatitis were estimated.

Two-sided p-values were provided with nominal level of statistical significance set to 5%. Analyses were performed using Stata.³⁶

Results

EPIC population characteristics. This study was based on a population of 476,106 participants, 70% women, with an overall median age at recruitment of 52 years. Within a

mean follow-up time of 14 years, and a total of 6,640,000 person-years, 1,283 incident pancreatic cancers were diagnosed (727 women) as reported in Table 1, with a median age at diagnosis of 67 years and age standardized incidence rate equal to 5.4 per 100,000 person-years.

Lifetime and baseline alcohol consumptions were 2- and 4-fold higher in men than in women, respectively. On average, beer and wine represented, respectively, 35% and 50% of total alcohol intake in men, and 12.5% and 63% in women. These patterns of consumption were consistent across countries in women, while consumptions were more heterogeneous in men. The proportion of non-drinkers was higher in women than in men. Men and women non-drinkers (< 0.1 g/day) differed by their educational attainment, physical activity level and diabetes mellitus status when they were compared to alcohol consumers. Percentage of smokers at recruitment was higher among alcohol drinkers than among alcohol non-drinkers. Characteristics by categories of baseline alcohol intake are shown into the Table 2.

Baseline alcohol intake. In men, baseline alcohol intake was statistically significantly associated with PC risk, with HR comparing alcohol intake greater than 60 g/day to the reference category (0.1-4.9g/day) equal to 1.63 (95%CI: 1.16, 2.29; $p_{\text{Wald}}=0.03$), as reported in Table 3. The association remained statistically significant when baseline alcohol intake was modelled as a continuous variable (HR for every increment of 12g/day: 1.05; 95% CI: 1.01, 1.09; $p_{\text{trend}}=0.02$). For women, no statistically significant association between baseline alcohol intake and PC risk was observed, either as a categorical ($p_{\text{Wald}}=0.68$) or as a continuous (HR for every increment of 12g/day: 1.04; 95% CI: 0.97, 1.12; $p_{\text{trend}}=0.28$) exposure.

Lifetime alcohol intake. Compared to the reference category, HR for men heavy drinkers (>60 g/day) was 1.77 (95% CI: 1.06, 2.95) without overall statistical significance among categories ($p_{\text{Wald}}=0.23$), as reported in Table 3. Analyses in continuous showed HR for a 12 g/day increase equal to 1.06 (95% CI: 1.02, 1.10; $p_{\text{trend}}<0.01$). No statistically significant associations were observed in women.

Type of alcoholic beverages. Mutually adjusted HR estimates for baseline alcoholic beverages are shown in Figure 1. Beer consumption was positively associated with PC risk with a 9% (95% CI: 1.02, 1.15; $p_{\text{trend}}=0.01$) and a 22% (95% CI: 1.03, 1.44; $p_{\text{trend}}=0.02$) risk increase for 12 g/day in men and women, respectively. The highest levels of beer consumption (>40 g/day in men and >20 g/day in women) were statistically significantly associated with PC risk compared to the reference category (0.1-2.9 g/day) with HR equal to 1.58 (95% CI: 1.10, 2.40) and 2.04 (95% CI: 1.13, 3.68) for men and women, respectively. Spirits/liquors in men

were associated with a 17% higher risk (95% CI: 1.04, 1.32; $p_{\text{trend}}=0.01$) for a 12 g/day increase, while no relationships were observed in women. Wine intake was not associated with PC risk, consistently in men and women. Similar results were observed for lifetime alcohol intake from the different beverages and PC risk (Supplementary Figure 1).

Dose-response relationship. Figure 2 illustrates the dose-response relationship of the baseline and lifetime alcohol intake and PC risk in men, using restricted cubic splines. The trend for baseline and lifetime alcohol intake suggests a linear-shaped association, without evidence for departure from linearity either for baseline ($p_{\text{non-linearity}}=0.83$) or lifetime alcohol ($p_{\text{non-linearity}}=0.57$).

Evaluating heterogeneity. Heterogeneity tests by sex and country for baseline alcohol intake were not statistically significant, with p-values equal to 0.63 (data not shown) and 0.33 (Supplementary Figure 2), respectively. Alcohol intake was not associated with PC risk among never smokers with HRs per 12g/day increase equal to 1.06 (95%CI: 0.98, 1.15; $p_{\text{trend}}=0.13$), unlike current smokers with HR equal to 1.05 (95%CI: 1.00, 1.11; $p_{\text{trend}}=0.04$). However, the overall interaction test for heterogeneity between alcohol and smoking status was not statistically significant ($p_{\text{heterog}}=0.84$) (Table 4). Thus, the association between baseline alcohol and PC risk was not different across smoking status.

Sensitivity analyses. After exclusion of the first two years of follow-up no substantial differences in results was observed in the association with baseline alcohol intake (data not shown). Among women, adjustment for menopausal status, ever use of hormone therapy, and number of full-term pregnancies in women did not alter estimates appreciably. The sensitivity analysis for external adjustment by history of chronic pancreatitis indicated that unadjusted HR estimate comparing baseline heavy drinkers (>60 g/day) vs. moderate drinkers (0.1-4.9 g/day) was marginally attenuated for estimates of relative risk between alcohol and chronic pancreatitis as large as 4 and estimates of the PC relative risk associated with chronic pancreatitis not exceeding 5. Larger attenuations of HR estimates were observed for more extreme scenarios, as displayed in Supplementary Table 1.

Discussion

In this study, alcohol was positively associated with PC risk in men, the relation being particularly apparent among heavy drinkers compared to light drinkers, consistently for baseline and lifetime alcohol intakes, controlling for a comprehensive list of confounding factors. There was no statistically significant association between alcohol consumption and

1 PC in women. Analyses by alcoholic subtypes showed positive relationships for beer and
2 spirits/liquors but not for wine. These results were virtually unaltered after sensitivity
3 analyses.

4 These findings support observations from other prospective studies.^{11,12,14,37,38} Our results
5 showed that each 12 g/day of alcohol in men was linearly associated with a 5% increase in PC
6 risk for baseline intakes, with a stronger association with the largest amounts of alcohol
7 greater than 60 g/day, consistently with results from the most recent meta-analyses.^{13,17,18}

8 While alcohol drinking has been related to PC risk in men, fewer studies found an association
9 in women.^{14,37} Women drink generally less than men,³⁹ as it was notably the case in the EPIC
10 study, the chance to observe a significant association with PC risk is weaker in women,
11 particularly if such association is apparent at high level of alcohol intake. However, no
12 evidence for heterogeneity across genders between alcohol and PC risk emerged in our study
13 ($p_{\text{heterog}}=0.63$), suggesting that an association with PC risk in women would have been
14 observed if they were showing exposure to alcohol as high as levels observed in men.

15 Our study used information on lifetime alcohol intake, less often investigated in relation to PC
16 risk. It revealed a statistically significant positive relationship with total lifetime alcohol
17 consumption in men, whether it was modelled as continuous variable with a 6% increase risk
18 for 12g/day or as categories, with men with the highest level of lifetime consumption
19 (>60g/day) having a 77% higher risk when compared to the light drinkers category. Although,
20 one case control study from California showed a more than three-fold significantly increased
21 OR for those with a history of binge drinking,⁴⁰ this association has not been shown in
22 previous prospective analyses.^{15,16,40}

23 Specific analyses on alcohol subtypes in this study showed that PC risk was statistically
24 significantly associated with spirits/liquors and beer in men, consistently using baseline and
25 lifetime intake. In women, results were more heterogeneous, showing associations with beer
26 intake at baseline, but not with lifetime intake. These findings are in line with previous studies
27 showing spirits/liquors consumption frequently associated with PC risk.^{12,14,16,18,37,38} However,
28 the association between beer consumption and PC risk was not reported in recent prospective
29 studies, especially in women. Our results also showed no association with wine intakes,
30 consistent observations with the other prospective studies.^{12,14,16,18,37} Moreover, country-
31 specific associations showed HR homogeneous estimates despite the variability of drinking
32 patterns across EPIC countries.

1 The consumption of alcoholic beverages leads to the production of acetaldehyde, the most
2 important metabolite derived from ethanol which increases the production of reactive oxygen
3 species and DNA-adducts.⁴¹ Acetaldehyde was classified as carcinogenic in 2012 by the
4 IARC Monograph program.¹⁰ Although oxidative stress produced by ethanol may induce
5 damage in pancreatic tissues through lipid peroxidation,^{42,43} associations observed in this
6 study varied depending on alcoholic subtypes. In vitro models investigating non-alcoholic
7 compounds of alcoholic beverages have shown that beer, unlike pure ethanol or wine, may
8 dose-dependently increase amylase secretion of rat's acinar cells, and potentially disturb
9 exocrine activity of the pancreas through alteration of cells' functions.⁴⁴ In parallel, the
10 absence of association between wine and PC risk could be partially explained by the fact that
11 wine contains molecules with anti-oxidative properties like polyphenols that may counteract
12 ethanol.⁴⁵ Resveratrol, a well-known polyphenolic compound of wine, has been reported to
13 suppress cell transformation, to induce apoptosis through a p53-dependent pathway and to
14 have chemo-preventive effects.⁴⁶ More recently, in vitro and ex-vivo models have shown
15 resveratrol suppressive action on pancreatic cells through inhibition of leukotriene A₄
16 hydrolase, an enzyme involved into pancreatic cancer cells growth.⁴⁷

17 It has been suggested that cigarette smoking in combination to ethanol may be associated with
18 pancreatic stellate cells activation in cells culture, which are the cells responsible for pancreas
19 fibrosis - a pre-cancerous lesion of PC.⁴⁸ Despite some evidence for interaction between
20 smoking and alcohol consumption on PC risk in case-control studies,¹⁹ this finding has not
21 been replicated in prospective studies,^{11,12,14} possibly due to the lack of sufficient statistical
22 power. In this study, no interaction between alcohol and smoking was observed, consistently
23 with one large American prospective study.¹⁴ This evidence lends further support to the
24 hypothesis that the relationship between alcohol and PC risk does not depend on smoking.

25 Our study has several strengths and limitations. We took advantage of the large number of PC
26 cases accrued in the EPIC study over a median of 14 years follow up, larger than previous
27 evaluations within EPIC,^{15,16} where no association was observed between alcohol intake and
28 PC. However, as EPIC participants are volunteers, they may be healthier and not
29 representative of the general population. Thus, the variability of alcohol intake could be lower
30 than in the general population. Moreover, self-reported assessments of alcohol intake are
31 prone to measurement errors, and could have biased the estimates of the association between
32 alcohol and PC risk. However, a previous calibration study in EPIC showed an absence of
33 impact in the assessment of the diet/disease association.²⁵

Study subjects with heavy alcohol consumption are susceptible to develop chronic pancreatitis,⁴⁹ a known risk factor for PC.⁵⁰ Accounting for chronic pancreatitis may provide useful information on the mechanism of the relationship between alcohol and PC risk. To address this, a sensitivity analysis was performed. For this analysis to be informative, a priori assumptions were set using evidence from the literature, i.e. the relative risk estimates of chronic pancreatitis associated with PC risk,³⁵ the prevalence of chronic pancreatitis among moderate drinkers³² and the relative risk estimates of chronic pancreatitis comparing extreme to light alcohol drinkers.³¹ The sensitivity analysis suggests that PC HR estimate in relation to alcohol intake was not substantially altered when information on chronic pancreatitis was accounted for, thus suggesting that alcohol intake exerts its carcinogenic role only partially through chronic pancreatitis.

Conclusion

In summary, our study has shown a moderate but statistically significant increase in PC risk with high alcohol intake, either baseline or lifetime, and particularly with beer and spirits/liquors. These findings provide epidemiologic evidence for the role of alcohol consumption as a potential carcinogen of the pancreas.

References

1. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015;136:E359-386.
2. Lepage C, Capocaccia R, Hackl M, Lemmens V, Molina E, Pierannunzio D, Sant M, Trama A, Faivre J. Survival in patients with primary liver cancer, gallbladder and extrahepatic biliary tract cancer and pancreatic cancer in Europe 1999–2007: Results of EUROCORE-5. *European Journal of Cancer* 2015;51:2169–78.
3. Rahib L, Smith BD, Aizenberg R, Rosenzweig AB, Fleshman JM, Matrisian LM. Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and pancreas cancers in the United States. *Cancer Res* 2014;74:2913–21.
4. Malvezzi M, Bertuccio P, Levi F, La Vecchia C, Negri E. European cancer mortality predictions for the year 2014. *Ann Oncol* 2014;25:1650–6.
5. Bray F, Ren J-S, Masuyer E, Ferlay J. Global estimates of cancer prevalence for 27 sites in the adult population in 2008. *Int J Cancer* 2013;132:1133–45.
6. American Institute for World Cancer Research Fund International. Pancreatic cancer | Continuous Update Project Report. Food, Nutrition, Physical Activity, and the Prevention of Pancreatic Cancer [Internet]. 2012;Available from: [http:// www.dietandcancerreport.org](http://www.dietandcancerreport.org)
7. Maisonneuve P, Lowenfels AB. Risk factors for pancreatic cancer: a summary review of meta-analytical studies. *Int J Epidemiol* 2015;44:186–98.
8. Barone E, Corrado A, Gemignani F, Landi S. Environmental risk factors for pancreatic cancer: an update. *Arch Toxicol* 2016;90:2617–42.
9. Ilic M, Ilic I. Epidemiology of pancreatic cancer. *World J Gastroenterol* 2016;22:9694–705.
10. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Personal habits and indoor combustions. Volume 100 E. A review of human carcinogens. *IARC Monogr Eval Carcinog Risks Hum* 2012;100:1–538.
11. Heinen MM, Verhage BAJ, Ambergen TAW, Goldbohm RA, van den Brandt PA. Alcohol consumption and risk of pancreatic cancer in the Netherlands cohort study. *Am J Epidemiol* 2009;169:1233–42.
12. Jiao L, Silverman DT, Schairer C, Thiébaud ACM, Hollenbeck AR, Leitzmann MF, Schatzkin A, Stolzenberg-Solomon RZ. Alcohol use and risk of pancreatic cancer: the NIH-AARP Diet and Health Study. *Am J Epidemiol* 2009;169:1043–51.
13. Tramacere I, Scotti L, Jenab M, Bagnardi V, Bellocchio R, Rota M, Corrao G, Bravi F, Boffetta P, La Vecchia C. Alcohol drinking and pancreatic cancer risk: a meta-analysis of the dose-risk relation. *Int J Cancer* 2010;126:1474–86.
14. Gapstur SM, Jacobs EJ, Deka A, McCullough ML, Patel AV, Thun MJ. Association of alcohol intake with pancreatic cancer mortality in never smokers. *Arch Intern Med* 2011;171:444–51.
15. Bueno de Mesquita HB, Maisonneuve P, Moerman CJ, Runia S, Boyle P. Lifetime consumption of alcoholic beverages, tea and coffee and exocrine carcinoma of the pancreas: a population-based case-control study in The Netherlands. *Int J Cancer* 1992;50:514–22.

16. Rohrmann S, Linseisen J, Vrieling A, Boffetta P, Stolzenberg-Solomon RZ, Lowenfels AB, Jensen MK, Overvad K, Olsen A, Tjønneland A, Boutron-Ruault M-C, Clavel-Chapelon F, et al. Ethanol intake and the risk of pancreatic cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Cancer Causes Control* 2009;20:785–94.
17. Bagnardi V, Rota M, Botteri E, Tramacere I, Islami F, Fedirko V, Scotti L, Jenab M, Turati F, Pasquali E, Pelucchi C, Galeone C, et al. Alcohol consumption and site-specific cancer risk: a comprehensive dose-response meta-analysis. *Br J Cancer* 2015;112:580–93.
18. Wang Y-T, Gou Y-W, Jin W-W, Xiao M, Fang H-Y. Association between alcohol intake and the risk of pancreatic cancer: a dose-response meta-analysis of cohort studies. *BMC Cancer* 2016;16:212.
19. La Torre G, Sferrazza A, Gualano MR, de Waure C, Clemente G, De Rose AM, Nicolotti N, Nuzzo G, Siliquini R, Boccia A, Ricciardi W. Investigating the Synergistic Interaction of Diabetes, Tobacco Smoking, Alcohol Consumption, and Hypercholesterolemia on the Risk of Pancreatic Cancer: A Case-Control Study in Italy. *Biomed Res Int* 2014;2014.
20. Riboli E, Hunt KJ, Slimani N, Ferrari P, Norat T, Fahey M, Charrondière UR, Hémond B, Casagrande C, Vignat J, Overvad K, Tjønneland A, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr* 2002;5:1113–24.
21. Kaaks R, Slimani N, Riboli E. Pilot phase studies on the accuracy of dietary intake measurements in the EPIC project: overall evaluation of results. European Prospective Investigation into Cancer and Nutrition. *Int J Epidemiol* 1997;26 Suppl 1:S26-36.
22. Slimani N, Ferrari P, Ocké M, Welch A, Boeing H, Liere M, Pala V, Amiano P, Lagiou A, Mattisson I, Stripp C, Engeset D, et al. Standardization of the 24-hour diet recall calibration method used in the European prospective investigation into cancer and nutrition (EPIC): general concepts and preliminary results. *Eur J Clin Nutr* 2000;54:900–17.
23. Sieri S, Agudo A, Kesse E, Klipstein-Grobusch K, San-José B, Welch AA, Krogh V, Luben R, Allen N, Overvad K, Tjønneland A, Clavel-Chapelon F, et al. Patterns of alcohol consumption in 10 European countries participating in the European Prospective Investigation into Cancer and Nutrition (EPIC) project. *Public Health Nutr* 2002;5:1287–96.
24. Slimani N, Deharveng G, Unwin I, Southgate D a. T, Vignat J, Skeie G, Salvini S, Parpinel M, Møller A, Ireland J, Becker W, Farran A, et al. The EPIC nutrient database project (ENDB): a first attempt to standardize nutrient databases across the 10 European countries participating in the EPIC study. *Eur J Clin Nutr* 2007;61:1037–56.
25. Ferrari P, Day NE, Boshuizen HC, Roddam A, Hoffmann K, Thiébaud A, Pera G, Overvad K, Lund E, Trichopoulou A, Tumino R, Gullberg B, et al. The evaluation of the diet/disease relation in the EPIC study: considerations for the calibration and the disease models. *Int J Epidemiol* 2008;37:368–78.
26. Thiébaud ACM, Bénichou J. Choice of time-scale in Cox's model analysis of epidemiologic cohort data: a simulation study. *Stat Med* 2004;23:3803–20.
27. Cust AE, Smith BJ, Chau J, van der Ploeg HP, Friedenreich CM, Armstrong BK, Bauman A. Validity and repeatability of the EPIC physical activity questionnaire: a validation study using accelerometers as an objective measure. *International Journal of Behavioral Nutrition and Physical Activity* 2008;5:33.

- 1 28. Schoenfeld D. Partial Residuals for The Proportional Hazards Regression Model.
2 *Biometrika* 1982;69:239–41.
- 3 29. Durrleman S, Simon R. Flexible regression models with cubic splines. *Stat Med*
4 1989;8:551–61.
- 5 30. Greenland S. Basic Methods for Sensitivity Analysis of Biases. *Int J Epidemiol*
6 1996;25:1107–16.
- 7 31. Samokhvalov AV, Rehm J, Roerecke M. Alcohol Consumption as a Risk Factor for
8 Acute and Chronic Pancreatitis: A Systematic Review and a Series of Meta-analyses.
9 *EBioMedicine* 2015;2:1996–2002.
- 10 32. Setiawan VW, Pandol SJ, Porcel J, Wilkens LR, Le Marchand L, Pike MC, Monroe
11 KR. Prospective Study of Alcohol Drinking, Smoking, and Pancreatitis: The Multiethnic
12 Cohort. *Pancreas* 2016;45:819–25.
- 13 33. Raimondi S, Lowenfels AB, Morselli-Labate AM, Maisonneuve P, Pezzilli R.
14 Pancreatic cancer in chronic pancreatitis; aetiology, incidence, and early detection. *Best Pract*
15 *Res Clin Gastroenterol* 2010;24:349–58.
- 16 34. Duell EJ, Lucenteforte E, Olson SH, Bracci PM, Li D, Risch HA, Silverman DT, Ji
17 BT, Gallinger S, Holly EA, Fontham EH, Maisonneuve P, et al. Pancreatitis and pancreatic
18 cancer risk: a pooled analysis in the International Pancreatic Cancer Case-Control Consortium
19 (PanC4). *Ann Oncol* 2012;mds140.
- 20 35. Dhar P, Kalghatgi S, Saraf V. Pancreatic Cancer in Chronic Pancreatitis. *Indian J Surg*
21 *Oncol* 2015;6:57–62.
- 22 36. StataCorp LP. Stata Statistical Software: Release 14. College Station, TX: StataCorp
23 LP. www.stata.com2015;
- 24 37. Genkinger JM, Spiegelman D, Anderson KE, Bergkvist L, Bernstein L, van den
25 Brandt PA, English DR, Freudenheim JL, Fuchs CS, Giles GG, Giovannucci E, Hankinson
26 SE, et al. Alcohol intake and pancreatic cancer risk: a pooled analysis of fourteen cohort
27 studies. *Cancer Epidemiol Biomarkers Prev* 2009;18:765–76.
- 28 38. Michaud DS, Vrieling A, Jiao L, Mendelsohn JB, Steplowski E, Lynch SM,
29 Wactawski-Wende J, Arslan AA, Bas Bueno-de-Mesquita H, Fuchs CS, Gross M, Helzlsouer
30 K, et al. Alcohol intake and pancreatic cancer: a pooled analysis from the pancreatic cancer
31 cohort consortium (PanScan). *Cancer Causes Control* 2010;21:1213–25.
- 32 39. Wilsnack RW, Wilsnack SC, Kristjanson AF, Vogeltanz-Holm ND, Gmel G. Gender
33 and alcohol consumption: patterns from the multinational GENACIS project. *Addiction*
34 2009;104:1487–500.
- 35 40. Gupta S, Wang F, Holly EA, Bracci PM. Risk of pancreatic cancer by alcohol dose,
36 duration, and pattern of consumption, including binge drinking: a population-based study.
37 *Cancer Causes Control* 2010;21:1047–59.
- 38 41. Seitz HK, Stickel F. Molecular mechanisms of alcohol-mediated carcinogenesis. *Nat*
39 *Rev Cancer* 2007;7:599–612.
- 40 42. Criddle DN, Raraty MGT, Neoptolemos JP, Tepikin AV, Petersen OH, Sutton R.
41 Ethanol toxicity in pancreatic acinar cells: mediation by nonoxidative fatty acid metabolites.
42 *Proc Natl Acad Sci USA* 2004;101:10738–43.
- 43 43. Palmieri VO, Grattagliano I, Palasciano G. Ethanol induces secretion of oxidized
44 proteins by pancreatic acinar cells. *Cell Biol Toxicol* 2007;23:459–64.

- 1 44. Gerloff A, Singer MV, Feick P. Beer and its Non-Alcoholic Compounds: Role in
2 Pancreatic Exocrine Secretion, Alcoholic Pancreatitis and Pancreatic Carcinoma. *Int J*
3 *Environ Res Public Health* 2010;7:1093–104.
- 4 45. Howard A, Chopra M, Thurnham DI, Strain JJ, Fuhrman B, Aviram M. Red wine
5 consumption and inhibition of LDL oxidation: what are the important components? *Medical*
6 *Hypotheses* 2002;59:101–4.
- 7 46. Jang M, Cai L, Udeani GO, Slowing KV, Thomas CF, Beecher CW, Fong HH,
8 Farnsworth NR, Kinghorn AD, Mehta RG, Moon RC, Pezzuto JM. Cancer chemopreventive
9 activity of resveratrol, a natural product derived from grapes. *Science* 1997;275:218–20.
- 10 47. Oi N, Jeong C-H, Nadas J, Cho Y-Y, Pugliese A, Bode AM, Dong Z. Resveratrol, a
11 Red Wine Polyphenol, Suppresses Pancreatic Cancer by Inhibiting Leukotriene A4
12 Hydrolase. *Cancer Res* 2010;70:9755–64.
- 13 48. Lee ATK, Xu Z, Pothula SP, Patel MB, Pirola RC, Wilson JS, Apte MV. Alcohol and
14 cigarette smoke components activate human pancreatic stellate cells: implications for the
15 progression of chronic pancreatitis. *Alcohol Clin Exp Res* 2015;39:2123–33.
- 16 49. Yadav D, Lowenfels AB. The Epidemiology of Pancreatitis and Pancreatic Cancer.
17 *Gastroenterology* 2013;144:1252–61.
- 18 50. Herreros-Villanueva M, Hijona E, Bañales JM, Cosme A, Bujanda L. Alcohol
19 consumption on pancreatic diseases. *World J Gastroenterol* 2013;19:638–47.

20

Financial disclosure

This work was supported by the Direction Générale de la Santé (French Ministry of Health) (Grant GR-IARC-2003-09-12-01), by the European Commission (Directorate General for Health and Consumer Affairs) and the International Agency for Research on Cancer. The national cohorts are supported by the Danish Cancer Society (Denmark); the Ligue Contre le Cancer, the Institut Gustave Roussy, the Mutuelle Générale de l'Éducation Nationale and the Institut National de la Santé et de la Recherche Médicale (France); the Deutsche Krebshilfe, the Deutsches Krebsforschungszentrum, and the Federal Ministry of Education and Research (Germany); the Hellenic Health Foundation, the Stavros Niarchos Foundation and the Hellenic Ministry of Health and Social Solidarity (Greece); the Italian Association for Research on Cancer and the National Research Council (Italy); the Dutch Ministry of Public Health, Welfare and Sports, the Netherlands Cancer Registry, LK Research Funds, Dutch Prevention Funds, the Dutch Zorg Onderzoek Nederland, the World Cancer Research Fund and Statistics Netherlands (the Netherlands); the Health Research Fund, Regional Governments of Andalucía, Asturias, Basque Country, Murcia (project 6236) and Navarra, Instituto de Salud Carlos III, Redes de Investigación Cooperativa (RD06/0020) (Spain); the Swedish Cancer Society, the Swedish Scientific Council and the Regional Government of Skåne (Sweden); Cancer Research UK (14136 to EPIC-Norfolk; C570/A16491 and C8221/A19170 to EPIC-Oxford), Medical Research Council (1000143 to EPIC-Norfolk, MR/M012190/1 to EPIC-Oxford) (United Kingdom), the Stroke Association, the British Heart Foundation, the Department of Health, the Food Standards Agency, and the Wellcome Trust (UK). This work was part of Sabine Naudin's PhD at Claude Bernard Lyon 1 University (France), funded by the Communauté de Recherche Académique de la Région Auvergne Rhône-Alpes (France).

Conflict of interest

None to declare.

Data sharing statement

For information on how to submit an application for gaining access to EPIC data and/or biospecimens, please follow the instructions at <http://epic.iarc.fr/access/index.ph>

Table 1. Country- and sex-specific frequencies of PC cases and other characteristics of the study population.

	Participants	PC cases	Follow-up (years) ¹	PY ²	Age (years) ³	Baseline alcohol (g/day) ⁴	BNC ²	Baseline beer (g/day) ⁴	Baseline wine (g/day) ⁴	Baseline spirits/liquors (g/day) ⁴	Lifetime alcohol (g/day) ⁴	% LNC ²
Men												
Italy	14,023	35	14	195,883	50	24 (1-54)	4	2 (0-3)	20 (0-47)	2 (0-5)	24 (2-49)	3
Spain	15,138	55	16	239,109	50	28 (0-67)	15	3 (0-9)	22 (0-56)	3 (0-11)	43 (4-91)	4
United Kingdoms	22,848	69	14	329,209	53	13 (0-34)	6	5 (0-22)	5 (0-12)	2 (0-3)	15 (1-34)	2
The Netherlands	9,627	20	15	140,422	44	18 (0-44)	9	10 (0-29)	4 (0-10)	4 (0-11)	⁻⁶	⁻⁶
Greece	10,814	25	10	112,787	52	18 (0-44)	10	4 (0-9)	9 (0-26)	5 (0-15)	30 (0-74)	6
Germany	21,177	72	10	219,536	53	24 (2-53)	4	15 (0-40)	8 (0-22)	2 (0-7)	27 (5-56)	1
Sweden	22,304	93	16	358,863	51	9 (0-23)	10	4 (0-8)	2 (0-8)	2 (0-7)	⁻⁶	⁻⁶
Denmark	26,287	187	15	382,609	56	28 (4-62)	2	14 (1-36)	10 (0-30)	3 (0-8)	21 (5-41)	1
All Men	142,218	556	14	1,978,417	53	20 (0-50)	7	7 (0-22)	10 (0-30)	3 (0-7)	26 (3-55)	2
Women												
France	67,395	56	13	869,319	52	11 (0-29)	14	1 (0-2)	9 (0-24)	0 (0-1)	7 (0-17)	15
Italy	30,511	69	14	434,978	51	9 (0-24)	23	1 (0-1)	7 (0-24)	0 (0-1)	6 (0-17)	15
Spain	24,848	51	16	398,811	48	4 (0-14)	53	1 (0-2)	3 (0-12)	0 (0-0)	4 (0-14)	39
United Kingdoms	52,56	119	15	793,462	48	7 (0-17)	6	1 (0-4)	5 (0-12)	1 (0-3)	9 (0-21)	4
The Netherlands	26,906	73	14	384,205	53	8 (0-24)	18	1 (0-2)	4 (0-11)	1 (0-3)	7 (0-18)	7
Greece	15,233	19	11	168,491	54	3 (0-9)	35	1 (0-1)	2 (0-6)	1 (0-2)	3 (0-10)	33
Germany	27,379	44	10	284,937	48	9 (0-23)	5	2 (0-7)	6 (0-17)	1 (0-1)	7 (1-15)	2
Sweden	26,368	111	17	442,242	51	5 (0-13)	18	1 (0-3)	2 (0-7)	0 (0-1)	⁻⁶	⁻⁶
Denmark	28,719	138	15	432,414	56	14 (1-34)	3	3 (0-7)	8 (0-30)	2 (0-5)	9 (1-19)	18
Norway	33,969	47	13	452,121	48	3 (0-7)	21	1 (0-2)	2 (0-5)	⁻⁵	⁻⁶	⁻⁶
All women	333,888	727	14	4,660,980	51	8 (0-22)	17	1 (0-4)	5 (0-15)	1 (0-2)	7 (0-17)	11
All participants	476,106	1283	14	6,639,397	52	12 (0-32)	14	3 (0-8)	7 (0-21)	1 (0-3)	13 (0-31)	8

¹ Means;

² PY: Person-years; BNC: Percentage of baseline non-consumers (<0.1 g/day); LNC: Percentage of lifetime never consumers (<0.1 g/day);

³ Medians;

⁴ Means (10th-90th percentiles);

⁵ Information on alcohol coming from spirits/liquors not available in Norway;

⁶ Information on lifetime alcohol not collected in Naples (Italy), Bilthoven (The Netherlands), Sweden, and Norway.

Table 2. Baseline characteristics of the EPIC participants by categories of baseline alcohol intake¹.

		Baseline alcohol intake categories (g/day)						
Men		Total	< 0.1	0.1 - 4.9	5 - 14.9	15 - 29.9	30 - 59.9	> 60
Number of participants	(n)	142,219	9,709	30,494	37,890	29,450	25,272	9,404
Number of PC cases	-	556	40	95	134	120	104	63
Person-Years	-	1,978,418	131,552	433,917	534,503	405,407	347,434	125,605
Age at recruitment	(years)	52 ± 10	54 ± 11	51 ± 12	52 ± 11	52 ± 9	53 ± 9	53 ± 8
Height	(cm)	175 ± 7	172 ± 8	175 ± 7	175 ± 7	175 ± 7	175 ± 7	174 ± 7
Weight	(kg)	81 ± 12	80 ± 13	80 ± 12	81 ± 12	81 ± 11	82 ± 12	83 ± 13
BMI	(kg/m ²)	27 ± 4	27 ± 4	26 ± 4	26 ± 4	26 ± 3	27 ± 4	27 ± 4
Diabetes mellitus	(%)	4	7	4	3	3	3	4
Smokers at recruitment	(%)	29	29	23	25	29	36	49
Participants moderately active	(%)	24	21	23	24	25	25	25
Participants with at least a university degree	(%)	27	15	23	29	31	29	22
Energy from non-alcohol drinking	(kcal/day)	2,268 ± 637	2,282 ± 686	2,222 ± 659	2,260 ± 626	2,297 ± 620	2,277 ± 620	2,318 ± 637
Lifetime alcohol intakes	(g/day)	26 ± 29	23 ± 47	10 ± 20	15 ± 15	24 ± 18	38 ± 23	65 ± 41
Baseline alcohol intakes	(g/day)	20 ± 23	0 ± 0	2 ± 1	9 ± 3	21 ± 4	42 ± 8	82 ± 25
Women	Units		< 0.1	0.1 - 4.9	5 - 14.9	15 - 29.9	> 30	
Number of participants	(n)	333,887	56,754	132,159	89,804	35,695	19,475	-
Number of PC cases	-	727	127	278	187	81	54	-
Person-Years	-	4,660,979	799,607	1,841,850	1,261,568	490,471	267,483	-
Age at recruitment	(years)	51 ± 10	52 ± 9	50 ± 10	51 ± 10	51 ± 9	52 ± 8	-
Height	(cm)	162 ± 7	160 ± 7	163 ± 7	163 ± 6	163 ± 6	163 ± 6	-
Weight	(kg)	66 ± 12	68 ± 13	66 ± 12	65 ± 11	64 ± 10	65 ± 11	-
BMI	(kg/m ²)	25 ± 4	26 ± 5	25 ± 5	24 ± 4	24 ± 4	24 ± 4	-
Diabetes melitus	(%)	2	4	2	1	1	2	-
Smokers at recruitment	(%)	19	17	18	18	22	31	-
Participants moderately active	(%)	27	22	29	28	27	26	-
Participants with at least a university degree	(%)	23	13	20	28	30	32	-
Energy from non-alcohol drinking	(kcal/day)	1,877 ± 529	1,831 ± 534	1,845 ± 521	1,910 ± 520	1,963 ± 540	1,915 ± 558	-
Lifetime alcohol intakes	(g/day)	7 ± 9	1 ± 6	3 ± 5	8 ± 7	14 ± 9	23 ± 14	-
Baseline alcohol intakes	(g/day)	8 ± 12	0 ± 0	2 ± 1	9 ± 3	21 ± 4	43 ± 15	-

¹ Means ± SD are presented for continuous variables, frequencies for categorical variables.

Table 3. Hazard Ratio (HR) estimates (95% CI) for baseline and lifetime alcohol intakes and PC.

Men		Baseline alcohol				Lifetime alcohol			
		cases	PY	HR ¹	95%CI	cases	PY	HR ²	95%CI
Continuous (12 g/day) ³		556	1,978,417	1.05	(1.01, 1.09)	429	1,460,432	1.06	(1.02, 1.10)
	P _{trend}				0.02				<0.01
Categories (g/day)									
	Ex-consumers	-	-	-	-	24	61,485	1.78	(0.75, 4.22)
	Non consumers	40	131,552	1.23	(0.84, 1.79)	4	33,366	0.53	(0.16, 1.74)
	0.1 - 4.9 ⁴	101	439,915	1.00	(Ref)	41	176,469	1.00	(Ref)
	5 - 14.9	132	532,427	0.99	(0.76, 1.29)	119	400,402	1.22	(0.82, 1.81)
	15 - 29.9	116	403,985	1.11	(0.83, 1.47)	116	389,206	1.26	(0.84, 1.90)
	30 - 59.9	104	345,443	1.10	(0.82, 1.47)	88	287,583	1.42	(0.93, 2.17)
	≥60 ⁵	63	125,095	1.63	(1.16, 2.29)	37	111,921	1.77	(1.06, 2.95)
	P _{Wald}				0.03				0.23
Women		Baseline alcohol				Lifetime alcohol			
		cases	PY	HR ¹	95%CI	cases	PY	HR ²	95%CI
Continuous (12 g/day) ³		727	4,660,980	1.04	(0.97, 1.12)	537	3,486,009	1.01	(0.88, 1.14)
	P _{trend}				0.28				0.90
Categories (g/day)									
	Ex-consumers	-	-	-	-	31	176,499	1.07	(0.54, 2.11)
	Non consumers	127	799,607	0.98	(0.78, 1.23)	63	495,243	0.72	(0.47, 1.10)
	0.1 - 4.9 ⁴	280	1,852,494	1.00	(Ref)	210	1,296,401	1.00	(Ref)
	5 - 14.9	187	1,257,465	1.00	(0.82, 1.21)	165	1,052,229	1.06	(0.85, 1.34)
	15 - 29.9	80	487,565	1.11	(0.86, 1.44)	59	382,037	1.16	(0.85, 1.59)
	≥30 ⁵	53	263,849	1.16	(0.85, 1.59)	9	83,601	0.93	(0.47, 1.85)
	P _{Wald}				0.68				0.79

¹ Models for baseline alcohol intake were stratified by center and age at recruitment. Systematic adjustment was undertaken for smoking intensity, physical activity level, educational attainment, diabetes status, BMI, height;


² Models for lifetime alcohol intake were stratified by center and age at recruitment. Systematic adjustment was undertaken for smoking intensity, physical activity level, educational attainment, diabetes status, BMI, height, duration of alcohol drinking, time since quitting, and an indicator variable for drinkers;

³ 12g of alcohol correspond to about one standard glass of either wine, beer or spirits;

⁴ The category 0.1 to 4.9 g/days was used as the reference category;

⁵ Wald test for overall significance, according to the χ^2 distribution with degrees of freedom equal to the number of categories minus one.

Figure 1. Baseline intake of beer, wine and spirits/liquors (g/day) and Hazard Ratio (HR) of pancreatic cancer in men and women.



¹ Models for baseline alcohol intake by subtypes were stratified by center and age at recruitment. Systematic adjustment was undertaken for smoking intensity, physical activity level, educational attainment, diabetes status, BMI, height, and baseline energy intake from other alcohol subtypes;

² p_{Wald} for overall significance across categories were performed according to the χ^2 distribution with degrees of freedom equal to the number of categories minus one. Trend tests were performed for continuous variable;

³ The category of light drinkers was used as the reference category (0.1-2.9 g/day for beer and wine, and 0.1-1.9 g/day for spirits/liquors);

⁴ 12g of alcohol correspond to about one standard glass of either wine, beer or spirits/liquors.

Table 4: Hazard Ratio¹ (95% CI) for overall pancreatic cancer risk by categories of baseline alcohol use (g/day) and smoking status (never and current smokers at baseline).

All participants	Never smokers				Current smokers				
	Cases	PY	HR ³	(95%CI)	Cases	PY	HR ³	(95%CI)	p _{heterogeneity} ⁴
Continuous (12 g/day) ²	494	3,286,210	1.06	(0.98, 1.15)	422	1,469,414	1.05	(1.00, 1.11)	
				0.13				0.04	0.84
P _{trend}									
Categories (g/day)									
Non consumers	86	578,541	0.94	(0.71, 1.23)	51	176026.5	1.26	(0.89, 1.77)	
0.1 - 4.9	182	1,261,449	1.00	(Ref)	105	439200.4	1.00	(Ref)	
5 - 14.9	126	879,891	0.99	(0.79, 1.25)	88	360975.2	0.91	(0.68, 1.22)	
15 - 29.9	60	359,073	1.08	(0.82, 1.50)	73	227177.8	1.19	(0.87, 1.63)	
30 - 59.9	28	176,457	0.93	(0.65, 1.48)	71	193786.8	1.25	(0.91, 1.73)	
≥ 60	12	30,799	2.17	(1.18, 3.99)	34	72247.58	1.50	(1.00, 2.28)	
p _{Wald} ⁵				0.14				0.10	0.41

¹ Models were stratified by centre, age at recruitment and sex. Systematic adjustment was undertaken for smoking status, physical activity level, educational attainment, diabetes status, BMI, height, and an indicator variable for drinkers;


² 12g of alcohol correspond to about one standard glass of either wine, beer or spirits/liquors;

³ Models included interaction terms between baseline alcohol use and a smoking indicator (0=never smokers; 1=current smokers), keeping as reference category the group of light alcohol users (0.1–4.9 g/day) among never smokers, whereas former smokers and participants without information on their smoking status were excluded;


⁴ Differences in HRs were assessed comparing the log-likelihood of models with and without interaction terms between alcohol and smoking status to one degree of freedom χ^2 distribution for analyses in continuous, and to five degrees of freedom χ^2 distribution for analyses in categories;

⁵ P-value was determined using a Wald test for contrasts according to a χ^2 distribution with five degrees of freedom.

Figure 2. Hazard Ratio (HR) functions and corresponding 95% confidence intervals (95% CI) describing the linear (dark blue) and the curvilinear (light blue) dose-response relationship between baseline and lifetime alcohol intake (g/day) and PC risk, according to pancreatic cancer frequencies in men.



Supplementary Figure 1. Lifetime intake of beer, wine and spirits/liquors (g/day) and Hazard Ratio (HR) of pancreatic cancer in men and women.




¹ Models for lifetime alcohol intake by subtypes were stratified by center and age at recruitment. Systematic adjustment was undertaken for smoking intensity, physical activity level, educational attainment, diabetes status, BMI, height, duration of subtype drinking, time since quitting subtype drinking, an indicator variable for drinkers of the subtype under analysis, and lifetime energy intake from other alcohol subtypes;

² Wald tests for overall significance across categories were performed according to the χ^2 distribution with degrees of freedom equal to the number of categories minus one. Trend tests were performed for continuous variable;

³ The category of light drinkers was used as the reference category (0.1-2.9 g/day for beer and wine, and 0.1-1.9 g/day for spirits/liquors);

⁴ 12g of alcohol correspond to about one standard glass of either wine, beer or spirits/liquors.

Supplementary Figure 2: Associations between alcohol intake per 12g/day increase and PC risk (and associated 95% CI) by country of origin, for men and women combined. P-value for heterogeneity was equal to 0.33, and was obtained by comparing the log-likelihood difference of models with and without interaction terms to a chi-square statistics with 9 degrees of freedom.



Supplementary Table 1. Sensitivity analysis for adjustment by history of chronic pancreatitis (Z) via external information in the relationship between baseline alcohol intake (X) and risk of pancreatic cancer (D).

p_{Z0} ¹	RR_{ZX} ²	RR_{DZ} ³ =	Estimates of HR_{DX} ⁴					
			1.5	2	2.5	5	10	15
0.005	2.0		1.64	1.63	1.63	1.61	1.57	1.54
0.01	2.0		1.63	1.62	1.62	1.58	1.52	1.46
0.02	2.0		1.62	1.61	1.60	1.53	1.43	1.36
0.005	4.0		1.63	1.62	1.60	1.55	1.46	1.38
0.01	4.0		1.62	1.59	1.57	1.48	1.32	1.21
0.02	4.0		1.60	1.56	1.52	1.36	1.15	1.02
0.005	6.0		1.62	1.60	1.58	1.50	1.36	1.24
0.01	6.0		1.60	1.57	1.53	1.39	1.18	1.04
0.02	6.0		1.57	1.51	1.45	1.23	0.98	0.83

¹ Prevalence of chronic pancreatitis among baseline moderate drinkers;

² Chronic pancreatitis RR estimates associated with baseline alcohol intake;

³ Pancreatic cancer RR estimates associated with chronic pancreatitis;

⁴ Estimates of pancreatic cancer Hazard Ratios associated with baseline alcohol intake adjusted for chronic pancreatitis. The pancreatic cancer HR estimate for heavy drinkers (>60g/day) compared to the reference category of moderate drinkers (0.1-4.9g/day) not adjusted for chronic pancreatitis was 1.64.